



Haycock, P. C., Burgess, S., Nounu, A., Zheng, J., Okoli, G. N., Bowden, J., Wade, K. H., Timpson, N. J., Evans, D. M., Willeit, P., Aviv, A., Gaunt, T. R., Hemani, G., Mangino, M., Ellis, H. P., Kurian, K. M., Pooley, K. A., Eeles, R. A., Lee, J. E. (2017). Association Between Telomere Length and Risk of Cancer and Non-Neoplastic Diseases: A Mendelian Randomization Study. *JAMA Oncology*, 3(5), 636-651. <https://doi.org/10.1001/jamaoncol.2016.5945>

Peer reviewed version

Link to published version (if available):

[10.1001/jamaoncol.2016.5945](https://doi.org/10.1001/jamaoncol.2016.5945)

[Link to publication record in Explore Bristol Research](#)

PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via JAMA at <http://jamanetwork.com/journals/jamaoncology/fullarticle/2604820>. Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

1 **Mendelian randomization study of the association between telomere length and risk of**  
2 **cancer and non-neoplastic diseases**

3

4 The Telomeres Mendelian Randomization Collaboration

5

6

7

8 Correspondence: Philip Haycock

9 MRC Integrative Epidemiology Unit

10 University of Bristol

11 Bristol

12 UK

13

14 [philip.haycock@bristol.ac.uk](mailto:philip.haycock@bristol.ac.uk)

15 Tel: +44 1173 310 088

16 2995 words [word limit 3000]

17 3 figures, 2 tables, 132 references; 7 supplementary figures / 6 supplementary tables

18

19

20

21

22

23

24

25

26 **ABSTRACT 349 WORDS**

27 **Importance** The causal direction and magnitude of the association between telomere length  
28 and incidence of cancer and non-neoplastic diseases is uncertain, due to the susceptibility of  
29 observational studies to confounding and reverse causation.

30 **Objective** To conduct a Mendelian randomization study, using germline genetic variants as  
31 instrumental variables, to appraise the causal relevance of telomere length for risk of cancer  
32 and non-neoplastic diseases.

33 **Data Sources** Genome-wide association studies (GWAS) published up to January 15 2015.

34 **Study Selection** GWAS of non-communicable diseases that assayed germline genetic  
35 variation and did not select cohort or control participants on the basis of pre-existing diseases.  
36 Of 163 GWAS of non-communicable diseases identified, summary data from 103 were  
37 available.

38 **Data Extraction** Summary association statistics for single nucleotide polymorphisms (SNPs)  
39 that are strongly associated with telomere length in the general population.

40 **Main Outcomes** Odds ratios (ORs) for disease per standard deviation (SD) higher telomere  
41 length due to germline genetic variation.

42 **Results** Summary data were available for 35 cancers and 48 non-neoplastic diseases,  
43 corresponding to 420,081 cases (median 2,526 per disease) and 1,093,105 controls (median  
44 6,789 per disease). Increased telomere length due to germline genetic variation was generally  
45 associated with increased risk for site-specific cancers. The strongest associations were  
46 observed for (ORs per 1-SD change in genetically increased telomere length): glioma 5.27  
47 (3.15-8.81), serous low-malignant-potential ovarian cancer 4.35 (2.39-7.94), lung  
48 adenocarcinoma 3.19 (2.40-4.22), neuroblastoma 2.98 (1.92-4.62), bladder cancer 2.19 (1.32-  
49 3.66), melanoma 1.87 (1.55-2.26), testicular cancer 1.76 (1.02-3.04), kidney cancer 1.55

50 (1.08-2.23) and endometrial cancer 1.31 (1.07-1.61). Associations were stronger for rarer  
51 cancers and at tissue sites with lower rates of stem cell division ( $P < 0.05$ ). There was  
52 generally little evidence of association between genetically increased telomere length and risk  
53 of psychiatric, autoimmune, inflammatory, diabetic and other non-neoplastic diseases, except  
54 for coronary heart disease (0.78 [0.67-0.90]), abdominal aortic aneurysm (0.63 [0.49-0.81]),  
55 celiac disease (0.42 [0.28-0.61]) and interstitial lung disease (0.09 [0.05- 0.15]).

56 **Conclusions** It is likely that longer telomeres increase risk for several cancers but reduce risk  
57 for some non-neoplastic diseases, including cardiovascular diseases.

58

59

60

61

62

63

64

65

66

67

68

69

70

## 71 INTRODUCTION

72

73 At the ends of chromosomes, telomeres are DNA-protein structures that protect the genome  
74 from damage, shorten progressively over time in most somatic tissues<sup>1</sup> and are proposed  
75 physiological markers of ageing.<sup>2,3</sup> Shorter leukocyte telomeres are correlated with older age,  
76 male sex and other known risk factors for non-communicable diseases<sup>4-6</sup> and are generally  
77 associated with higher risk for cardiovascular diseases<sup>7,8</sup>, type 2 diabetes<sup>9</sup> and non-vascular  
78 non-neoplastic causes of mortality.<sup>8</sup> Whether these associations are causal, however, is  
79 unknown. Telomere length has also been implicated in risk of cancer but the direction and  
80 magnitude of the association is uncertain and contradictory across observational studies.<sup>10-14</sup>  
81 The uncertainty reflects the considerable difficulty of designing observational studies of  
82 telomere length and cancer incidence that are robust to reverse causation, confounding and  
83 measurement error.

84 The aim of the present report was to conduct a Mendelian randomization study, using  
85 germline genetic variants as instrumental variables for telomere length, to help clarify the  
86 nature of the association between telomere length and risk of cancer and non-neoplastic  
87 diseases. The approach, which mimics the random allocation of individuals to the placebo  
88 and intervention arms of a randomized controlled trial, allowed us to: (1) estimate the  
89 direction and broad magnitude of the association of telomere length with risk of multiple  
90 cancer and non-neoplastic diseases; (2) appraise the evidence for causality in the estimated  
91 etiological associations; (3) investigate potential sources of heterogeneity in findings for site-  
92 specific cancers; and (4) compare genetic estimates to findings based on directly measured  
93 telomere length in prospective observational studies.

94

## 95    **METHODS**

96

### 97    *Study design*

98    The design of our study, illustrated in Figure S1, had three key components: 1) the  
99    identification of genetic variants to serve as instruments for telomere length; 2) the  
100    acquisition of summary data for the genetic instruments from genome wide association  
101    studies (GWASs) of diseases and risk factors for non-communicable diseases; and 3) the  
102    classification of diseases and risk factors into primary or secondary outcomes based on *a*  
103    *priori* statistical power. As a first step, we searched the GWAS catalog<sup>15,16</sup> on the 15 January  
104    2015, to identify single nucleotide polymorphisms (SNPs) associated with telomere length.  
105    To supplement the list with additional potential instruments, we also searched the original  
106    study reports curated by the GWAS catalog (using a P-value threshold of  $5 \times 10^{-8}$ ).<sup>17–25</sup> We  
107    acquired summary data for all SNPs identified by our search from a meta-analysis of GWASs  
108    of telomere length, involving 9,190 participants of European ancestry.<sup>18</sup>

109    The second key component of our design strategy involved the acquisition of summary data,  
110    corresponding to the selected genetic instruments for telomere length, from GWASs of non-  
111    communicable diseases and risk factors (Fig. S1). As part of this step, we invited principal  
112    investigators of non-communicable disease studies curated by the GWAS catalog<sup>15,26</sup> to share  
113    summary data for our study. We also downloaded summary data for diseases and risk factors  
114    from publically available sources, including study-specific websites, dbGAP, ImmunoBase  
115    and the GWAS catalog (Fig. S1).

116    The third key component of our design strategy was the classification of diseases and risk  
117    factors into either primary or secondary outcomes, which we defined on the basis of *a priori*  
118    statistical power to detect associations with telomere length. Primary outcomes were defined

as diseases with sufficient cases and controls for >50% statistical power and secondary outcomes defined as diseases with <50% statistical power to detect odds ratios  $\geq 2.0$  per standard deviation (SD) change in genetically increased telomere length (alpha assumed to be 0.01). All risk factors were defined as secondary outcomes. Risk factors with <50% statistical power were excluded.

Further details on our design strategy can be found in the supplement.

### *Comparison with prospective observational studies*

We searched PubMed for prospective observational studies of the association between telomere length and disease (see Tables S3 and S4 for details of the search strategy and inclusion criteria). Study-specific relative risks for disease per unit change or quantile comparison of telomere length were transformed to a SD scale using previously described methods.<sup>27</sup> Hazard ratios, risk ratios and odds ratios were assumed to approximate the same measure of relative risk. Where multiple independent studies of the same disease were identified, these were combined by fixed effects meta-analysis, unless there was strong evidence of between-study heterogeneity ( $P_{\text{Cochran's } Q} < 0.001$ ), in which case they were kept separate.

### *Statistical analysis*

We combined summary data across SNPs into a single instrument, using maximum likelihood to estimate the slope of the relationship between  $\beta_{\text{GD}}$  and  $\beta_{\text{GP}}$  and a variance-covariance matrix to make allowance for linkage disequilibrium between SNPs,<sup>28</sup> where  $\beta_{\text{GD}}$  is the change in disease log odds or risk factor levels per copy of the effect allele and  $\beta_{\text{GP}}$  is the SD change in telomere length per copy of the effect allele (see supplementary methods

for technical details). The slope from this approach can be interpreted as the log odds ratio for binary outcomes, or the unit change for continuous risk factors, per SD change in genetically increased telomere length. P-values for heterogeneity amongst SNPs, in the estimated associations of genetically increased telomere length with disease and risk factors, were estimated by likelihood ratio tests.<sup>28</sup> Associations between genetically increased telomere length and continuous risk factors were transformed into SD units. For five secondary disease outcomes where only a single SNP was available for analysis, we estimated associations using the Wald ratio:  $\beta_{GD}/\beta_{GP}$ , with standard errors approximated by the delta method.<sup>29</sup>

Inference of causality in the estimated etiological associations between telomere length and disease depends on satisfaction of Mendelian randomization assumptions (Fig. S7; see Table S6 for a glossary of terms).<sup>30,31</sup> The assumptions are: 1) the selected SNPs are associated with telomere length; 2) the selected SNPs are not associated with confounders; and 3) the selected SNPs are associated with disease exclusively through their effect on telomere length. If these assumptions are satisfied, the selected SNPs are valid instrumental variables and their association with disease can be interpreted as a causal effect of telomere length. We modeled the impact of violations of these assumptions through two sets of sensitivity analyses: a weighted median function<sup>32</sup> and MR-Egger regression<sup>30</sup> (see supplementary methods for technical details). We restricted our sensitivity analyses to diseases showing the strongest evidence of association with genetically increased telomere length (defined as  $P_{\text{Bonferroni}} \leq 0.05$ ).

We used meta-regression to appraise potential sources of heterogeneity in our findings for cancer. The association of genetically increased telomere length with the log odds of cancer was regressed on cancer incidence, survival time and median age-at-diagnosis, downloaded from the National Cancer Institute Surveillance, Epidemiology, and End Results (SEER)



Program,<sup>33</sup> and tissue-specific rates of stem cell division from Tomasetti and Vogelstein.<sup>34</sup> As the downloaded cancer characteristics from SEER correspond to the United States population, 77% of which was of white ancestry in 2015<sup>35</sup>, the meta-regression analyses excluded genetic studies conducted in East Asian populations.

All analyses were performed in R version 3.1.2<sup>36</sup> and Stata release 13.1 (StataCorp, College Station, TX). P-values were two-sided and evidence of association was declared at  $P < 0.05$ . Where indicated, Bonferroni corrections were used to make allowance for multiple testing, although this is likely to be overly conservative given the non-independence of many of the outcomes tested.

## RESULTS

We selected 16 SNPs as instruments for telomere length (Fig. S1 & Table 1). The selected SNPs correspond to 10 independent genomic regions that collectively account for 2-3% of the variance in leukocyte telomere length, which is equivalent to an F statistic of ~18. This indicates that the genetic instrument, constructed from these 10 independent genomic regions, is strongly associated with telomere length (details in supplementary discussion).<sup>37</sup> Summary data for the genetic instruments were available for 83 non-communicable diseases, corresponding to 420,081 cases (median 2,526 per disease) and 1,093,105 controls (median 6,789 per disease), and 44 risk factors (Fig. S1, Table 2 and Table S1). The median number of SNPs available across diseases was 11 (min=1, max=13) and across risk factors was 12 (min=11, max=13). Of the 83 diseases, 56 were classified as primary outcomes and 27 as secondary outcomes (Table 2, Fig. S1 and Table S1). For 9 of the 83 non-communicable diseases, additional summary data were available from 10 independent studies for replication

analyses, corresponding to 40,465 cases (median 1,416 per disease) and 52,306 controls (median 3,537 per disease) (Table S1).

The results from primary analyses of non-communicable diseases are presented in Figure 1; results from secondary analyses of risk factors and diseases with low *a priori* power are presented in the supplement (Fig. S2, S5 and S6). Genetically increased telomere length was associated with higher odds of disease for 9 of 22 primary cancers ( $P < 0.05$ ), including (odds ratio [95% confidence interval]): glioma (5.27 [3.15-8.81]), endometrial cancer (1.31 [1.07-1.61]), kidney cancer (1.55 [1.08-2.23]), testicular germ cell cancer (1.76 [1.02-3.04]), melanoma (1.87 [1.55-2.26]), bladder cancer (2.19 [1.32-3.66]), neuroblastoma (2.98 [1.92-4.62]), lung adenocarcinoma (3.19 [2.40-4.22]) and serous low-malignancy-potential (LMP) ovarian cancer (4.35 [2.39-7.94]) (Fig. 1). The associations were, however, highly variable across cancer types, varying from an odds ratio of 0.86 (0.50-1.48) for head and neck cancer to 5.27 (3.15-8.81) for glioma. Substantial variability was also observed within tissue sites. For example, the odds ratio for lung adenocarcinoma was 3.19 (2.40-4.22) compared to 1.07 (0.82-1.39) for squamous cell lung cancer. For serous LMP ovarian cancer the odds ratio was 4.35 (2.39-7.94) compared to odds ratios of 1.21 (0.87-1.68) for endometrioid ovarian cancer, 1.12 (0.94-1.34) for serous invasive ovarian cancer, 1.04 (0.66-1.63) for clear cell ovarian cancer and 1.04 (0.73-1.47) for mucinous ovarian cancer. The strongest evidence of association was observed for glioma, lung adenocarcinoma, neuroblastoma and serous LMP ovarian cancer ( $P_{\text{Bonferroni}} < 0.05$ ). Results for glioma and bladder cancer showed evidence for replication in independent datasets (independent datasets were not available for other cancers) (Fig. S3).

Genetically increased telomere length was associated with reduced odds of disease for 6 of 32 primary non-neoplastic diseases ( $P < 0.05$ ), including coronary heart disease (0.78 [0.67-0.9]), abdominal aortic aneurysm (0.63 [0.49-0.81]), Alzheimer's disease (0.84 [0.71-0.98]), celiac

disease (0.42 [0.28-0.61]), interstitial lung disease (0.09 [0.05-0.15]) and type 1 diabetes (0.71 [0.51-0.98]) ( $P < 0.05$ ) (Figure 1). The strongest evidence of association was observed for coronary heart disease ( $P_{\text{Bonferroni}} = 0.05$ ) and abdominal aortic aneurysm, celiac disease and interstitial lung disease ( $P_{\text{Bonferroni}} < 0.05$ ). The associations with coronary heart disease and interstitial lung disease showed evidence for replication in independent datasets (Fig. S3).

Our genetic findings were generally similar in direction and magnitude to estimates based on observational prospective studies of leukocyte telomere length and disease (Figure 3). Our genetic estimates for lung adenocarcinoma, melanoma, kidney cancer and glioma, were, however, stronger in comparison to observational estimates.

In sensitivity analyses, we appraised the potential impact of confounding by pleiotropic pathways on our results. Associations estimated by the weighted median and MR-Egger were broadly similar to the main results for glioma, lung adenocarcinoma, serous LMP ovarian cancer, neuroblastoma, abdominal aortic aneurysm, coronary heart disease and interstitial lung disease (Fig. S4). In the second set of sensitivity analyses, implemented by MR-Egger regression, we found little evidence for the presence of pleiotropy ( $P_{\text{intercept}} \geq 0.27$ ) (Fig. S4). The MR-Egger analyses were, however, generally underpowered, as reflected by the wide confidence intervals in the estimated odds ratios.

In meta-regression analyses, we observed that genetically increased telomere length tended to be more strongly associated with rarer cancers ( $P = 0.02$ ) and cancers at tissue-sites with lower rates of stem cell division ( $P = 0.02$ ) (Figure 2). The associations showed little evidence of varying by percentage survival five years after diagnosis or median age-at-diagnosis ( $P \geq 37$ ).

## DISCUSSION

In this report we show that genetically increased telomere length is associated with increased risk of several cancers and with reduced risk of some non-neoplastic diseases. Given the random distribution of genotypes in the general population with respect to lifestyle and other environmental factors, as well as the fixed nature of germline genotypes, these results should be less susceptible to confounding and reverse causation in comparison to observational studies. Our results could, however, reflect violations of Mendelian randomization assumptions, such as confounding by pleiotropy, population stratification or ancestry.<sup>38</sup> Although we cannot entirely rule out this possibility, the majority of our results persisted in sensitivity analyses that made allowance for violations of Mendelian randomization assumptions. Confounding by population stratification or ancestry is also unlikely, given the adjustments made for ancestry in the original disease GWASs (see supplementary discussion). Our results are therefore compatible with causality.

### *Comparison with previous studies*

Our findings for cancer are generally contradictory to those based on retrospective studies, which tend to report increased risk for cancer in individuals with shorter telomeres.<sup>11,12,39–42</sup> The contradictory findings may reflect reverse causation in the retrospective studies, whereby shorter telomeres arise as a result of disease, or of confounding effects, e.g. due to cases being slightly older than controls even in age-matched analyses. Our findings for cancer are generally more consistent with those based on prospective observational studies, which tend to report weak or null associations of longer leukocyte telomeres with overall and site-specific risk of cancer,<sup>10–13,41,43–62</sup> with some exceptions.<sup>63</sup> Our results are also similar to previously reported Mendelian randomization studies of telomere length and risk of

melanoma, lung cancer, chronic lymphocytic leukemia and glioma.<sup>64-67</sup> The shape of the association with cancer may not, however, be linear over the entire telomere length distribution. For example, individuals with dyskeratosis congenita, a disease caused by germline loss-of-function mutations in the telomerase component genes *TERC* and *TERT*, have chronically short telomeres and are at increased risk of some cancers, particularly acute myeloid leukemia and squamous cell carcinomas arising at sites of leukoplakia,<sup>68,69</sup> presumably due to increased susceptibility to genome instability and chromosomal end-to-end fusions.<sup>70</sup> Our results should therefore be interpreted as reflecting the average association at the population level and may not be generalizable to the extreme ends of the telomere length distribution.

#### *Mechanisms of association*

Our cancer findings are compatible with known biology.<sup>70</sup> By limiting the proliferative potential of cells, telomere shortening may serve as a tumour suppressor; and individuals with longer telomeres may be more likely to acquire somatic mutations owing to increased proliferative potential.<sup>70</sup> Rates of cell division are, however, highly variable amongst tissues<sup>34</sup> and thus the relative gain in cell proliferative potential, conferred by having longer telomeres, may also be highly variable across tissues. This could explain the ~6-fold variation in odds ratios observed across cancer types in the present study, as well as the tendency of our results to be stronger at tissue sites with lower rates of stem cell division. For example, the association was strongest for glioma (OR=5.27) and comparatively weak for colorectal cancer (OR=1.09) and the rates of stem cell division in the tissues giving rise to these cancers differ by several orders of magnitude. In neural stem cells, which give rise to gliomas, the number of divisions is ~270 million and for colorectal stem cells is ~1.2 trillion over the average lifetime of an individual.<sup>34</sup> The observation that genetically increased telomere

length was more strongly associated with rarer cancers potentially reflects the same mechanism, since rarer cancers also tend to show lower rates of stem cell division.<sup>34</sup> For example, the incidence of glioma is 0.4 and for colorectal cancer is 42.4 per 100,000 per year in the United States.<sup>33</sup>

The inverse associations observed for some non-neoplastic diseases may reflect the impact of telomere shortening on tissue degeneration and an evolutionary trade-off for greater resistance to cancer at the cost of greater susceptibility to degenerative diseases, particularly cardiovascular diseases.<sup>71,72</sup>

### *Study limitations*

Our study is subject to some limitations, in addition to the Mendelian randomization assumptions already considered above. First, our method assumes that the magnitude of the association between SNPs and telomere length is consistent across tissues. Second, our study assumed a linear shape of association between telomere length and disease risk, whereas the shape could be “J” or “U” shaped.<sup>44,57,68</sup> Third, our results assume that the samples used to define the genetic instrument for telomere length<sup>18</sup> and the various samples used to estimate the SNP-disease associations are representative of the same general population, practically defined as being of similar ethnicity, age and sex distribution.<sup>73</sup> This assumption would, for example, not apply in the case of the SNP-disease associations derived from East Asian or pediatric populations. Generally speaking, violation of the aforementioned assumptions could bias the magnitude of the association between genetically increased telomere length and disease; but would be unlikely to increase the likelihood of false positives (i.e. incorrectly inferring an association when none exists).<sup>74</sup> Our results should therefore remain informative for the direction and broad magnitude of the average association at the population level, even

in the presence of such violations. Fourth, we cannot rule out chance in explaining some of the weaker findings. Fifth, our results may not be fully representative of non-communicable diseases (since not all studies shared data and our analyses were underpowered for the secondary disease outcomes). The diseases represented in our primary analyses probably account for >60% of all causes of death in American adults.<sup>75</sup>

#### *Clinical relevance of findings*

Our findings suggest that potential clinical applications of telomere length, e.g. as a tool for risk prediction or as an intervention target for disease prevention, may have to consider a trade-off in risk between cancer and non-neoplastic diseases. For example, a number of companies have been established that offer telomere length measurement services to the public (via a requesting physician), under the claim that shorter telomeres are a general indicator of poorer health status and older biological age and that such information can be used to motivate healthy lifestyle choices in individuals. However, the conflicting direction of association between telomere length and risk of cancer and non-neoplastic diseases, indicated by our findings, suggests that such services to the general public may be premature.

#### *Conclusion*

It is likely that longer telomeres increase risk for several cancers but reduce risk for some non-neoplastic diseases, including cardiovascular diseases. Further research is required to resolve whether telomere length is a useful predictor of risk that can help guide therapeutic interventions, to clarify the shape of any dose-response relationships and to characterise the nature of the association in population subgroups.

# The Telomeres Mendelian Randomization Collaboration

Philip Haycock<sup>1</sup>; Stephen Burgess<sup>2</sup>; Aayah Nounu<sup>1</sup>; Jie Zheng<sup>1</sup>; George N Okoli<sup>3</sup>; Jack Bowden<sup>1,4</sup>; Kaitlin Wade<sup>1</sup>; Nicholas Timpson<sup>1</sup>; David M. Evans<sup>1,5</sup>; Peter Willeit<sup>2,6</sup>; Abraham Aviv<sup>7</sup>; Tom R. Gaunt<sup>1</sup>; Gibran Hemani<sup>1</sup>; Massimo Mangino<sup>8,9</sup>; Hayley Patricia Ellis<sup>10</sup>; Kathreena Mary Kurian<sup>10</sup>; Karen Pooley<sup>11</sup> on behalf of the BCAC and OCAC consortia; Rosalind Eeles<sup>12</sup> on behalf of the PRACTICAL consortium; Jeffrey E Lee<sup>13</sup>; Shenying Fang<sup>13</sup>; Wei Chen<sup>13</sup>; Matthew H Law<sup>14</sup>; Lisa M Bowdler<sup>15</sup> and Mark M Iles<sup>16</sup> on behalf of the Melanoma meta-analysis consortium; Qiong Yang<sup>17</sup>; Bradford B. Worrall<sup>18</sup> and Hugh Stephen Markus<sup>19</sup> on behalf of the METASTROKE project of the ISGC; Rayjean J. Hung<sup>20,21</sup> and Chris I Amos<sup>22</sup> on behalf of the ILCCO consortium; Amanda Spurdle<sup>23</sup>; Deborah J Thompson<sup>24</sup> and Tracy O'Mara<sup>23</sup> on behalf of the ECAC consortium; Brian Wolpin<sup>25</sup>; Laufey Amundadottir<sup>26</sup> and Rachael Stolzenberg-Solomon<sup>27</sup> on behalf of the PanScan consortium; Antonia Trichopoulou<sup>29,30</sup>; Charlotte Onland-Moret<sup>31</sup>; Eiliv Lund<sup>32</sup>; Eric J Duell<sup>33</sup>; Federico Canzian<sup>34</sup>; Gianluca Severi<sup>35,36,37,38</sup>; Kim Overvad<sup>39</sup>; Marc J Gunter<sup>40</sup>; Rosario Tumino<sup>41</sup> and Ulrika Svenson<sup>42</sup> on behalf of EPIC; Andre van Rij<sup>43</sup>; Annette F Baas<sup>44</sup>; Matthew J Bown<sup>45</sup>; Nilesh J Samani<sup>45</sup>; Paul IW de Bakker<sup>44</sup>; Femke NG van t'Hof<sup>44</sup>; Gerard Tromp<sup>46,47</sup>; Gregory T Jones<sup>43</sup>; Helena Kuivaniemi<sup>46,47</sup> and James R Elmore<sup>48</sup> on behalf of the Aneurysm Consortium; Mattias Johansson<sup>49</sup>; James Mckay<sup>49</sup>; Ghislaine Scelo<sup>49</sup>; Robert Carreras-Torres<sup>49</sup>; Valerie Gaborieau<sup>49</sup>; Paul Brennan<sup>49</sup>; Paige M. Bracci<sup>50</sup>; Rachel E Neale<sup>15</sup>; Sara H Olson<sup>51</sup>; Steven Gallinger<sup>20</sup>; Donghui Li<sup>52</sup>; Gloria M. Petersen<sup>54</sup>; Harvey Risch<sup>55</sup>; and Alison P. Klein<sup>56</sup> on behalf of PanC<sup>4</sup>; Jiali Han<sup>57,58</sup>; Christian C. Abnet<sup>59</sup>; Neal D. Freedman<sup>59</sup>; Philip R. Taylor<sup>59</sup>; John M Maris<sup>60</sup>; Katja K Aben<sup>61,62</sup>; Lambertus A Kiemeny<sup>61</sup>; Sita H Vermeulen<sup>61</sup>; John K Wiencke<sup>63,64</sup>; Kyle M Walsh<sup>63,64</sup>; Margaret Wrensch<sup>63,64</sup>; Terri Rice<sup>63</sup>; Clare Turnbull<sup>65</sup>; Kevin Litchfield<sup>66</sup>; Lavinia Paternoster<sup>1</sup> and Marie Standl<sup>67</sup> on behalf of the EAGLE consortium; Gonçalo R Abecasis<sup>68</sup>; John Paul SanGiovanni<sup>69</sup>; Yong Li<sup>70</sup> and Vladan Mijatovic<sup>71</sup> on behalf of the CKDGen consortium; Yadav Sapkota<sup>15</sup>; Siew-Kee Low<sup>72</sup>; Krina T Zondervan<sup>73,74</sup>; Grant W Montgomery<sup>15</sup>; Dale R. Nyholt<sup>75,15</sup>; David A van Heel<sup>76</sup>; Karen Hunt<sup>76</sup>; Dan E. Arking<sup>77</sup>; Foram N. Ashar<sup>77</sup> and Nona Sotoodehnia<sup>78</sup> on behalf of the CHARGE-Sudden Cardiac Arrest Working Group; Daniel Woo<sup>79</sup>; Jonathan Rosand<sup>80</sup>; Mary Comeau<sup>81</sup>; W. Mark Brown<sup>82</sup>; Edwin K. Silverman<sup>83</sup>; John E Hokanson<sup>84</sup> and Michael Cho<sup>83</sup> on behalf of COPDGene; Jennie Hui<sup>85,86,87,88</sup>; Manuel Ferreira<sup>15</sup> and Philip J. Thompson<sup>89</sup> on behalf of the AAGC consortium; Alanna C. Morrison<sup>90</sup>; Janine F Felix<sup>91</sup> and Nicholas L



375 Smith<sup>92</sup> on behalf of the CHARGE-Heart Failure Working Group; Angela M Christiano<sup>93</sup>;  
 376 Lynn Petukhova<sup>94</sup>; Regina C. Betz<sup>95</sup>; Xing Fan<sup>96</sup>; Xuejun Zhang<sup>96</sup>; Caihong Zhu<sup>96</sup>; Carl  
 377 Langefeld<sup>97</sup>; Susan D. Thompson<sup>98</sup>; Feijie Wang<sup>99</sup>; Xu Lin<sup>99,100</sup>; David A. Schwartz<sup>101</sup>; Tasha  
 378 Fingerlin<sup>102</sup>; Jerome I. Rotter<sup>103,104</sup>; Mary Frances Cotch<sup>105</sup> and Richard A Jensen on behalf of  
 379 the CHARGE-Eye Working Group<sup>106,107</sup>; Matthias Munz<sup>108</sup>; Henrik Dommisch<sup>108</sup> and Arne  
 380 S Schaefer<sup>108</sup> on behalf of the European Periodontitis Genetics Group; Fang Han<sup>109</sup>; Hanna M  
 381 Ollila<sup>110</sup>; Ryan P. Hillary<sup>110</sup>; Omar Albagha<sup>111</sup>; Stuart H. Ralston<sup>112</sup>; Chenjie Zeng<sup>113</sup>; Wei  
 382 Zheng<sup>113</sup>; Xiao-Ou Shu<sup>113</sup>; Andre Reis<sup>114</sup>; Steffen Uebe<sup>114</sup>; Ulrike Hüffmeier<sup>114</sup>; Yoshiya  
 383 Kawamura<sup>115</sup>; Takeshi Otowa<sup>116,117</sup> and Tsukasa Sasaki<sup>118</sup> on behalf of the Japanese  
 384 Collaboration Team for GWAS of Panic Disorder; Martin Lloyd Hibberd<sup>119</sup>; Michael  
 385 Levin<sup>120</sup>; Sonia Davila<sup>121</sup>; Gang Xie<sup>122,20</sup>; Katherine Siminovitch<sup>122,20</sup>; Jin-Xin Bei<sup>123</sup>; Yi-Xin  
 386 Zeng<sup>123,124</sup>; Asta Försti<sup>125,126</sup>; Bowang Chen<sup>125</sup>; Stefano Landi<sup>127</sup>; Andre Franke<sup>128</sup>; Annegret  
 387 Fischer<sup>128,129</sup>; David Ellinghaus<sup>128</sup>; Carlos Flores<sup>130,131</sup>; Imre Noth<sup>132</sup>; Shwu-Fan Ma<sup>132</sup>; Jia  
 388 Nee Foo<sup>133</sup>; Jianjun Liu<sup>133</sup>; Jong-Won Kim<sup>134</sup>; David G. Cox<sup>135</sup>; Olivier Delattre<sup>136</sup>; Olivier  
 389 Mirabeau<sup>136</sup>; Christine F. Skibola<sup>137</sup>; Clara S. Tang<sup>138</sup>; Merce Garcia-Barcelo<sup>138</sup>; Kai-Ping  
 390 Chang<sup>139</sup>; Wen-Hui Su<sup>140</sup>; Yu-Sun Chang<sup>141</sup>; Nicholas G Martin<sup>15</sup>; Scott Gordon<sup>15</sup>; Tracey  
 391 Wade<sup>142</sup>; Chaeyoung Lee<sup>143</sup>; Michiaki Kubo<sup>144</sup>; Pei-Chieng Cha<sup>145</sup>; Yusuke Nakamura<sup>146</sup>;  
 392 Daniel Levy<sup>147</sup>; Masayuki Kimura<sup>7</sup>; Shih-Jen Hwang<sup>147</sup>; Steven Hunt<sup>148</sup>; Tim Spector<sup>8</sup>;  
 393 Nicole Soranzo<sup>149</sup>; Ani W Manichaikul<sup>150</sup>; R Graham Barr<sup>151</sup>; Bratati Kahali<sup>152</sup>; Elizabeth  
 394 Speliotes<sup>152</sup> and Laura M Yerges-Armstrong<sup>153</sup> on behalf of the GOLD Consortium; Ching-  
 395 Yu Cheng<sup>154,155,156</sup>; Jost B. Jonas<sup>157,158</sup> and Tien Yin Wong<sup>154,155,156</sup> on behalf of the SEED  
 396 consortium; Isabella Fogh<sup>159</sup>; Kuang Lin<sup>159</sup> and John F. Powell<sup>159</sup> on behalf of the SLAGEN  
 397 and ALSGEN consortia; Kenneth Rice<sup>160</sup> on behalf of the ICBP; Caroline Relton<sup>1</sup>; Richard  
 398 M Martin<sup>1,3,161</sup>; George Davey Smith<sup>1</sup>

- 1 MRC Integrative Epidemiology Unit, University of Bristol, Bristol, UK
- 2 Department of Public Health and Primary Care, University of Cambridge
- 3 School of Social and Community Medicine, University of Bristol, Bristol, UK
- 4 MRC Biostatistics Unit, Cambridge, UK.
- 5 University of Queensland Diamantina Institute, Translational Research Institute, Brisbane, Queensland, Australia
- 6 Department of Neurology, Innsbruck Medical University, Austria
- 7 Center of Human Development and Aging, Department of Pediatrics, New Jersey Medical School, Rutgers, The State University of New Jersey
- 8 Department of Twin Research and Genetic Epidemiology, King's College London, London UK
- 9 NIHR Biomedical Research Centre at Guy's and St. Thomas' Foundation Trust, London, UK
- 10 Brain Tumour Research Group, Institute of Clinical Neuroscience, Learning and Research Building, Southmead Hospital, University of Bristol
- 11 Strangeways Research Laboratory, University of Cambridge, Cambridge, UK
- 12 The Institute of Cancer Research, London, UK
- 13 Department of Surgical Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX.
- 14 Statistical Genetics, QIMR Berghofer Medical Research Institute, Brisbane, Australia
- 15 QIMR Berghofer Medical Research Institute, Brisbane, Australia
- 16 Section of Epidemiology and Biostatistics, Leeds Institute of Cancer and Pathology, University of Leeds, Leeds, UK
- 17 Department of Biostatistics, Boston University School of Public Health, Boston, Massachusetts, United States of America and the Framingham Heart Study, Framingham, Massachusetts, United States of America
- 18 Departments of Neurology and Public Health Sciences University of Virginia Charlottesville, Virginia 22908
- 19 Department of Clinical Neurosciences, University of Cambridge, UK
- 20 Lunenfeld-Tanenbaum Research Institute of Mount Sinai Hospital, Toronto, Ontario, Canada
- 21 Division of Epidemiology, Dalla Lana School of Public Health, University of Toronto, 60 Murray St. Rm L5-215, Box 18, Toronto, ON M5T 3L9, Canada
- 22 Geisel School of Medicine, Dartmouth College
- 23 Genetics and Computational Biology Division, QIMR Berghofer Medical Research Institute, Brisbane, QLD 4006, Australia
- 24 Centre for Cancer Genetic Epidemiology, Department of Public Health and Primary Care, University of Cambridge, Cambridge, CB1 8RN, UK.
- 25 Dana-Farber Cancer Institute
- 26 Laboratory of Translational Genomics, Division of Cancer Epidemiology & Genetics, National Cancer Institute
- 27 Division of Cancer Epidemiology and Genetics, National Cancer Institute, NIH, DHHS.
- 29 Hellenic Health Foundation, Athens, Greece
- 30 WHO Collaborating Center for Nutrition and Health, Unit of Nutritional Epidemiology and Nutrition in Public Health, Dept. of Hygiene, Epidemiology and Medical Statistics, University of Athens Medical School, Greece
- 31 Dept of Epidemiology, Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht
- 32 Institute of community medicine, UiT The Arctic University of Norway
- 33 Unit of Nutrition and Cancer, Cancer Epidemiology Research Program, Bellvitge Biomedical Research Institute (IDIBELL), Catalan Institute of Oncology (ICO), Avda Gran Via 199-203, 08908 L'Hospitalet de Llobregat, Barcelona, Spain
- 34 Genomic Epidemiology Group, German Cancer Research Center (DKFZ), Heidelberg, Germany
- 35 Université Paris-Saclay, Université Paris-Sud, UVSQ, CESP, INSERM, Villejuif, France
- 36 Gustave Roussy, F-94805, Villejuif, France
- 37 Human Genetics Foundation (HuGeF), Torino, Italy

38 Cancer Council Victoria and University of Melbourne, Australia

39 Department of Public Health, Section for Epidemiology, Aarhus University, Aarhus, Denmark

40 School of Public Health, Imperial College London, London W2 1PG

41 Cancer Registry, Azienda Ospedaliera "Civile M.P.Arezzo", via Dante 109, Ragusa, IT

42 Department of Medical Biosciences, Umea University, Umea, Sweden

43 Surgery Department, University of Otago, Dunedin, New Zealand

44 Department of Medical Genetics, Center for Molecular Medicine and Department of Epidemiology,  
Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, The  
Netherlands

45 The Department of Cardiovascular Sciences and the NIHR Leicester Cardiovascular Biomedical  
Research Unit, University of Leicester, Leicester, LE2 7LX, UK.

46 Division of Molecular Biology and Human Genetics, Department of Biomedical Sciences, Faculty of  
Medicine and Health Sciences, Stellenbosch University, Tygerberg, 7505, South Africa

47 The Sigfried and Janet Weis Center for Research, Geisinger Health System, Danville, PA, USA

48 Department of Vascular and Endovascular Surgery, Geisinger Health System, Danville, PA 17822,  
USA

49 Genetic Epidemiology Group, International Agency for Research on Cancer, Lyon, France

50 Department of Epidemiology and Biostatistics, University of California San Francisco, San Francisco,  
California

51 Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York,  
New York, USA.

52 Department of Gastrointestinal Medical Oncology, University of Texas MD Anderson Cancer Center,  
Houston, Texas, USA

54 Department of Health Sciences Research, Mayo Clinic College of Medicine, Rochester, Minnesota,  
USA

55 Yale School of Public Health & Yale School of Medicine & Yale Cancer Center, 60 College St., PO  
Box 208034, New Haven, CT 06520-8034

56 Departments of Oncology, Pathology and Epidemiology, Johns Hopkins School of Medicine, Baltimore  
Maryland 21231

57 Department of Epidemiology, Fairbanks School of Public Health, Indiana University

58 Indiana University Melvin and Bren Simon Cancer Center, Indianapolis, IN, USA

59 Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville MD USA

60 Children's Hospital of Philadelphia, Perelman School of Medicine at the University of Pennsylvania

61 Radboud university medical center, Radboud Institute for Health Sciences, Nijmegen, The Netherlands

62 Netherlands Comprehensive Cancer Organization, Utrecht, The Netherlands

63 Department of Neurological Surgery, University of California, San Francisco, San Francisco, CA

64 Institute of Human Genetics, University of California, San Francisco, San Francisco, CA

65 William Harvey Research Institute, Queen Mary University, London, UK

66 Division of Genetics and Epidemiology, The Institute of Cancer Research, London, UK

67 Institute of Epidemiology I, Helmholtz Zentrum München - German Research Center for  
Environmental Health, Neuherberg, Germany

69 National Eye Institute, Division of Epidemiology and Clinical Research Clinical Trials Branch National  
Institutes of Health, Bethesda, USA

70 Department of Internal Medicine IV, University Hospital Freiburg

71 Department of Life and Reproduction Sciences, University of Verona

72 Laboratory of Statistical Analysis, Centre for Integrative Medical Sciences, The Institute of Physical  
and Chemical Research (RIKEN), Yokohama, Japan

73 Genetic and Genomic Epidemiology Unit, Wellcome Trust Centre for Human Genetics, University of  
Oxford, Oxford, UK

74 Nuffield Department of Obstetrics and Gynecology, University of Oxford, John Radcliffe Hospital,  
Oxford, UK

75 Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane,  
Australia

76 Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of  
London, London E1 2AT, UK

77 McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University School of Medicine,  
Baltimore, MD USA

78 Division of Cardiology and Cardiovascular Health Research Unit, Department of Medicine, University  
of Washington, Seattle, Washington 98101

79 University of Cincinnati College of Medicine, Department of Neurology, Cincinnati, OH, USA 45267

80 Massachusetts General Hospital, Neurology, Center for Human Genetic Research, MA, USA

81 Center for Public Health Genomics, Department of Biostatistical Sciences, Division of Public Health  
Sciences, Wake Forest School of Medicine, Medical Center Blvd, Winston-Salem, NC 27157

82 Department of Biostatistical Sciences, Division of Public Health Sciences, Wake Forest School of  
Medicine, Medical Center Blvd, Winston-Salem, NC 27157

83 Channing Division of Network Medicine, Brigham and Women's Hospital, Boston, MA 02115

83 Channing Division of Network Medicine, Brigham and Women's Hospital, Boston, MA, 02115

84 Department of Epidemiology, University of Colorado Anschutz Medical Campus, Aurora, Colorado,  
USA

85 Busselton Population Medical Research Institute Inc, Sir Charles Gairdner Hospital, Perth, Australia

86 PathWest Laboratory Medicine of Western Australia (WA), Perth, Australia

87 School of Pathology and Laboratory Medicine, University of WA, Perth, Australia

88 School of Population Health, University of WA, Perth, Australia

89 The Lung Health Clinic and Institute for Respiratory Health, University of Western Australia, Perth,  
Australia

90 Department of Epidemiology, Human Genetics, and Environmental Sciences, University of Texas  
Health Science Center at Houston, Houston, TX 77030 USA

91 Department of Epidemiology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, the  
Netherlands

92 Department of Epidemiology, University of Washington, Seattle WA 98101 USA

93 Departments of Dermatology and Genetics & Development, Columbia University, New York, NY, US

94 Departments of Dermatology and Epidemiology, Columbia University, New York, NY, US

95 Institute of Human Genetics, University of Bonn, Bonn D-53127, Germany

96 Institute of Dermatology & Department of Dermatology, First Affiliated Hospital of Anhui Medical  
University

97 Director, Center for Public Health Genomics, Department of Biostatistical Sciences, Division of Public  
Health Sciences, Wake Forest School of Medicine, Medical Center Blvd, Winston-Salem, NC 27157

98 Center for Autoimmune Genomics and Etiology, Cincinnati Children's Hospital Medical Center,  
Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH, USA

99 Institute for Nutritional Sciences, SIBS, Chinese Academy of Sciences, Shanghai, 200031, PR China

100 Key Laboratory of Nutrition and Metabolism, Chinese Academy of Sciences, Shanghai, 200031, PR  
China

101 University of Colorado, 12631 East 17th Avenue, B178, Aurora, CO 80045

102 Department of Biomedical Research, National Jewish Health Hospital

103 Institute for Translational Genomics and Population Sciences, Los Angeles Biomedical Research  
Institute at Harbor-UCLA Medical Center

104 Departments of Pediatrics and Medicine, 1124 W. Carson Street, Harbor-UCLA Medical Center,  
Torrance, CA 90502

105 Epidemiology Branch, Division of Epidemiology and Clinical Applications, NIH Intramural Research  
Program, National Eye Institute, National Institutes of Health, Clinical Research Center 3A2521,

106 Cardiovascular Health Research Unit, University of Washington, Seattle, Washington, USA  
 107 Department of Medicine, University of Washington, Seattle, Washington, USA  
 108 Charité – University Medicine Berlin, CC 03, Institute of Dental, Oral and Maxillary Medicine, Dept.  
 of Periodontology and Synoptic Dentistry, Aßmannshauser Str. 4-6, 14197 Berlin, Germany  
 109 Department of Pulmonary Medicine, Peking University People's Hospital, 100044 Beijing, China  
 110 Stanford University, Center for Sleep Sciences, Palo Alto, CA, USA  
 111 Centre for Genomic and Experimental Medicine, Institute of Genetics and Molecular Medicine,  
 University of Edinburgh, Western General Hospital, Edinburgh EH4 2XU, UK  
 112 Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, UK  
 113 Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Vanderbilt-  
 Ingram Cancer Center, Vanderbilt University Medical Center, Nashville, Tennessee  
 114 Institute of Human Genetics, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, Germany  
 115 Department of Psychiatry, Shonan Kamakura General Hospital, Kanagawa, Japan  
 116 Department of Neuropsychiatry, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan  
 117 Graduate School of Clinical Psychology, Teikyo Heisei University Major of Professional Clinical  
 Psychology, Tokyo, Japan  
 118 Department of Physical and Health Education, Graduate School of Education, The University of Tokyo,  
 Tokyo, Japan  
 119 Infectious Diseases, Genome Institute of Singapore, Singapore  
 120 Division of Infectious diseases, Department of medicine, Imperial College London, UK  
 121 Human genetics, Genome Institute of Singapore, Singapore  
 122 Departments of Medicine, Immunology, Molecular Genetics, University of Toronto  
 123 Sun Yat-sen University Cancer Center, State Key Laboratory of Oncology in South China,  
 Collaborative Innovation Center for Cancer Medicine Guangzhou 510060, P. R. China  
 124 Peking Union Medical College, Beijing 100730, P.R. China  
 125 Molecular Genetic Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany  
 126 Center for Primary Health Care Research, Clinical Research Center, Lund University, Malmö, Sweden  
 127 Department of Biology, University of Pisa, Pisa, Italy  
 128 Institute of Clinical Molecular Biology, Christian-Albrechts-University of Kiel, Kiel, Germany.  
 129 University Hospital Schleswig-Holstein, Kiel, Germany  
 130 Research Unit, Hospital Universitario N.S. de Candelaria, Tenerife, Spain  
 131 CIBER de Enfermedades Respiratorias, Instituto de Salud Carlos III, Madrid, Spain  
 132 Section of Pulmonary and Critical Care Medicine, University of Chicago, 5841 S. Maryland Ave.,  
 Chicago IL 60637-6076  
 133 Human Genetics, Genome Institute of Singapore, A\*STAR, Singapore 138672, Singapore  
 134 Dept. of Laboratory Medicine and Genetics, Samsung Medical Center, Sungkyunkwan, University  
 School of Medicine, Ilwon-dong 50, Gangnam-gu, Seoul, Korea, 135-710  
 135 Cancer Research Center of Lyon, INSERM U1052, Lyon, France  
 136 Inserm U830, Institut Curie, PSL University, 26 rue d'Ulm, 75248 Paris Cedex 05 France, France.  
 137 Department of Epidemiology, University of Alabama at Birmingham. 1665 University Boulevard,  
 Birmingham, AL 35294-0022, USA  
 138 Department of Surgery, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong,  
 SAR China  
 139 Department of Otolaryngology - Head and Neck Surgery, Chang Gung Memorial Hospital at Lin-Kou,  
 Taoyuan, Taiwan,  
 140 Department of Biomedical Sciences, Graduate Institute of Biomedical Sciences, College of Medicine,  
 Molecular Medicine Research Center, Chang Gung University, Taoyuan, Taiwan  
 141 Molecular Medicine Research Center, Chang Gung University, Taoyuan, Taiwan

- 142 School of Psychology, Flinders University
- 143 School of Systems Biomedical Science, Soongsil University, 369 Sangdo-ro, Dongjak-gu, Seoul 156-743, Korea
- 144 RIKEN Center for Integrative Medical Science, 1-7-22, Suehiro-cho, Tsurumi-ku, Yokohama, Kanagawa 230-0045, JAPAN
- 145 Division of Molecular Brain Science, Kobe University Graduate School of Medicine, 7-5-1 Kusunoki-chou, Chuo-ku, Kobe 650-0017, Japan
- 146 Center for Personalized Therapeutics, The University of Chicago, 900E 57th Street, Chicago IL 60637 USA
- 147 The NHLBI's Framingham Heart Study, Framingham, MA, Population Sciences Branch of the National Heart, Lung, and Blood Institute, Bethesda, MD.
- 148 Department of Genetic Medicine, Weill Cornell Medicine in Qatar, Doha, Qatar
- 149 Human Genetics, Wellcome Trust Sanger Institute, Genome Campus, Hinxton Cambridge
- 150 Center for Public Health Genomics, Department of Public Health Sciences, University of Virginia, Charlottesville, VA USA
- 151 Department of Medicine and Department of Epidemiology, Columbia University Medical Center, New York, NY 10032, USA
- 152 Department of Internal Medicine, Division of Gastroenterology and Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI 48109, USA
- 153 Department of Medicine, University of Maryland, Baltimore, MD 21201, USA
- 154 Singapore Eye Research Institute, Singapore National Eye Center, Singapore 168751, Singapore
- 155 Department of Ophthalmology, National University of Singapore and National University Health System, Singapore
- 156 Duke-National University of Singapore Graduate Medical School, Singapore
- 157 Beijing Institute of Ophthalmology, Beijing Tongren Eye Center, Beijing Tongren Hospital, Capital Medical University, Beijing Ophthalmology and Visual Science Key Lab, Beijing, China
- 158 Department of Ophthalmology, Medical Faculty Mannheim of the Ruprecht-Karls-University Heidelberg, Mannheim, Germany
- 159 Department of Basic and Clinical Neuroscience, Maurice Wohl Clinical Neuroscience Institute, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom
- 160 Dept of Biostatistics, University of Washington, Seattle, USA
- 161 University of Bristol / University Hospitals Bristol NHS Foundation Trust National Institute for Health Research Bristol Nutrition Biomedical Research Unit, Bristol, UK

399

400 Affiliations of the Telomeres Mendelian Randomization Collaboration

401

402 **Acknowledgements**

**Access to Data Statement:** Dr Philip C Haycock had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

**Role of the Funder/Sponsor:** The funding institutions had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

This work was supported by CRUK grant number C18281/A19169 (the Integrative Cancer Epidemiology Programme). Dr Haycock is supported by CRUK Population Research Postdoctoral Fellowship C52724/A20138. The MRC Integrative Epidemiology Unit is supported by grants MC\_UU\_12013/1 and MC\_UU\_12013/2. Dr Martin is supported by the National Institute for Health Research (NIHR), the Bristol Nutritional Biomedical Research Unit and the University of Bristol.

We gratefully acknowledge all the studies and databases that made GWAS summary data available (see supplementary materials for detailed acknowledgements): **AC** (the aneurysm consortium), **ALSGEN** (the International Consortium on Amyotrophic Lateral Sclerosis Genetics), **AMD Gene** (Age-related Macular Degeneration Gene Consortium), **BCAC** (Breast Cancer Association Consortium), **C4D** (Coronary Artery Disease Genetics Consortium), **CARDIoGRAM** (Coronary ARtery Disease Genome wide Replication and Meta-analysis), **CHARGE-HF** (Cohorts for Heart and Aging Research in Genomic Epidemiology Consortium – Heart Failure Working Group), **COPDGene** (The Genetic Epidemiology of Chronic Obstructive Pulmonary Disease), **CORECT** (ColoRectal Transdisciplinary Study), **CKDGen** (Chronic Kidney Disease Genetics consortium), **dbGAP** (database of Genotypes and Phenotypes), **DIAGRAM** (DIAbetes Genetics Replication And Meta-analysis), **EAGLE** (EARly Genetics & Lifecourse Epidemiology Eczema Consortium, excluding 23andMe), **ECAC** (Endometrial Cancer Association Consortium), **EGG** (Early Growth Genetics Consortium), **EPG** (European Periodontitis Genetics Group), **GABRIEL** (A Multidisciplinary Study to Identify the Genetic and Environmental Causes of Asthma in the European Community), **GCAN** (Genetic Consortium for Anorexia Nervosa), **GECCO** (Genetics and Epidemiology of Colorectal Cancer Consortium), **GIANT** (Genetic Investigation of ANthropometric Traits), **GLGC** (Global Lipids Genetics Consortium),

428 **GUGC** (Global Urate and Gout consortium), **ICBP** (International Consortium for Blood  
 429 Pressure), **IGAP** (International Genomics of Alzheimer's Project), **HPFS** (Health  
 430 Professionals Follow-Up Study), **JCTGPD** (Japanese Collaboration Team for GWAS of  
 431 Panic Disorder), **ILCCO** (International Lung Cancer Consortium), **ImmunoBase** (genetic  
 432 database of immunologically related human diseases), **IMSGC** (International Multiple  
 433 Sclerosis Genetic Consortium), **IIBDGC** (International Inflammatory Bowel Disease  
 434 Genetics Consortium); **KIDRISK** (Kidney cancer consortium), **MAGIC** (Meta-Analyses of  
 435 Glucose and Insulin-related traits Consortium), **MC** (the melanoma meta-analysis  
 436 consortium), **MESA** (Multi-Ethnic Study of Atherosclerosis), **METASTROKE/ISGC**  
 437 (METASTROKE project of the International Stroke Genetics Consortium), **NBCS** (Nijmegen  
 438 Bladder Cancer Study), **NHGRI-EBI GWAS catalog** (National Human Genome Research  
 439 Institute and European Bioinformatics Institute Catalog of published genome-wide  
 440 association studies), **NHS** (Nurses' Health Study), **OCAC** (Ovarian Cancer Association  
 441 Consortium), **PanScan** (Pancreatic Cancer Cohort Consortium), **PGC** (Psychiatric Genomics  
 442 Consortium), **PRACTICAL** (Prostate Cancer Association Group to Investigate Cancer  
 443 Associated Alterations in the Genome), **SEEDS** (the Singapore Epidemiology of Eye  
 444 Diseases Study), **SLAGEN** (Italian Consortium for the Genetics of Amyotrophic Lateral  
 445 Sclerosis), **SSGAC** (Social Science Genetics Association Consortium), **TAG** (Tobacco and  
 446 Genetics Consortium), **T1Dbase** (type 1 diabetes database), **TICG** (Tourette International  
 447 Collaborative-Genetics); **TSAICG** (Tourette Syndrome Association International Consortium  
 448 for Genetics).

449 We gratefully acknowledge the assistance and contributions of Dr Julia Gumy, Ms Lisa  
 450 Wright, Dr Georg B. Ehret (ICBP), Dr Louise V. Wain (ICBP), Dr Caroline Fox (CKDGen),  
 451 Dr Stephan Ripke (IIBDGC), Dr Jimmy Liu (IIBDGC), Dr Carl Anderson (IIBDGC), Dr



452 Jeremiah Scharf (TSAICG and TICG), Dr Lars Fritsche (AMD Gene), Dr Joanne Elena and  
453 Dr Paul KH Tam (Hirschsprung's disease GWAS).

**Table 1.** Single nucleotide polymorphisms associated with telomere length

| SNPs       | Chr | Pos       | Gene          | EA | OA | EAF* | Beta* | SE*    | P-value* | Phet* | No. studies* | Sample size* | Discovery p-value | % variance explained | Discovery study            |
|------------|-----|-----------|---------------|----|----|------|-------|--------|----------|-------|--------------|--------------|-------------------|----------------------|----------------------------|
| rs11125529 | 2   | 54248729  | <i>ACYP2</i>  | A  | C  | 0.16 | 0.065 | 0.012  | 0.000606 | 0.313 | 6            | 9177         | 8.00E-10          | 0.080                | Codd <sup>21</sup>         |
| rs6772228  | 3   | 58390292  | <i>PXK</i>    | T  | A  | 0.87 | 0.041 | 0.014  | 0.049721 | 0.77  | 6            | 8630         | 3.91E-10          | 0.200                | Pooley <sup>17</sup>       |
| rs12696304 | 3   | 169763483 | <i>TERC</i>   | C  | G  | 0.74 | 0.090 | 0.011  | 5.41E-08 | 0.651 | 6            | 9012         | 4.00E-14          | 0.319                | Codd <sup>22</sup>         |
| rs10936599 | 3   | 169774313 | <i>TERC</i>   | C  | T  | 0.76 | 0.100 | 0.011  | 1.76E-09 | 0.087 | 6            | 9190         | 3.00E-31          | 0.319                | Codd <sup>21</sup>         |
| rs1317082  | 3   | 169779797 | <i>TERC</i>   | A  | G  | 0.71 | 0.097 | 0.011  | 4.57E-09 | 0.029 | 6            | 9176         | 1.00E-08          | 0.319                | Mangino <sup>18</sup>      |
| rs10936601 | 3   | 169810661 | <i>TERC</i>   | C  | T  | 0.74 | 0.087 | 0.011  | 8.64E-08 | 0.433 | 6            | 9150         | 4.00E-15          | 0.319                | Pooley <sup>17</sup>       |
| rs7675998  | 4   | 163086668 | <i>NAF1</i>   | G  | A  | 0.80 | 0.048 | 0.012  | 0.008912 | 0.077 | 6            | 9161         | 4.35E-16          | 0.190                | Codd <sup>21</sup>         |
| rs2736100  | 5   | 1286401   | <i>TERT</i>   | C  | A  | 0.52 | 0.085 | 0.013  | 2.14E-05 | 0.54  | 4            | 5756         | 4.38E-19          | 0.310                | Codd <sup>21</sup>         |
| rs9419958  | 10  | 103916188 | <i>OBFC1</i>  | T  | C  | 0.13 | 0.129 | 0.013  | 5.26E-11 | 0.028 | 6            | 9190         | 9.00E-11          | 0.171                | Mangino <sup>18</sup>      |
| rs9420907  | 10  | 103916707 | <i>OBFC1</i>  | C  | A  | 0.14 | 0.142 | 0.014  | 1.14E-11 | 0.181 | 6            | 9190         | 7.00E-11          | 0.171                | Codd <sup>21</sup>         |
| rs4387287  | 10  | 103918139 | <i>OBFC1</i>  | A  | C  | 0.14 | 0.120 | 0.013  | 1.40E-09 | 0.044 | 6            | 8541         | 2.00E-11          | 0.171                | Levy <sup>25</sup>         |
| rs3027234  | 17  | 8232774   | <i>CTC1</i>   | C  | T  | 0.83 | 0.103 | 0.012  | 2.75E-08 | 0.266 | 6            | 9108         | 2.00E-08          | 0.292                | Mangino <sup>18</sup>      |
| rs8105767  | 19  | 22032639  | <i>ZNF208</i> | G  | A  | 0.25 | 0.064 | 0.011  | 0.000169 | 0.412 | 6            | 9096         | 1.11E-09          | 0.090                | Codd <sup>21</sup>         |
| rs412658   | 19  | 22176638  | <i>ZNF676</i> | T  | C  | 0.35 | 0.086 | 0.010  | 1.83E-08 | 0.568 | 6            | 9156         | 1.00E-08          | 0.484                | Mangino <sup>18</sup>      |
| rs6028466  | 20  | 39500359  | <i>DHX35</i>  | A  | G  | 0.17 | 0.058 | 0.013  | 0.003972 | 0.533 | 6            | 9190         | 2.57E-08†         | 0.041                | Mangino <sup>18</sup> & Gu |
| rs755017   | 20  | 63790269  | <i>ZBTB46</i> | G  | A  | 0.17 | 0.019 | 0.0129 | 0.339611 | 0.757 | 5            | 8026         | 6.71E-09          | 0.090                | Codd <sup>21</sup>         |

\*Summary data from Mangino et al<sup>18</sup>; Chr, chromosome; pos, base-pair position (GRCh38.p3); EA, effect allele, OA, other allele, Beta, standard deviation change in telomere length per copy of the effect allele; SE, standard error; EAF - effect allele frequency; Phet - p value for between-study heterogeneity in association between SNP and telomere length; †from a meta-analysis of Mangino<sup>18</sup> and Gu<sup>20</sup> performed in the present study.

**Table 2.** Study characteristics for primary non-communicable diseases

|   | No.<br>cases | No.<br>controls | No.<br>SNPs | Statistical<br>power | Pop. | Study / First author                          |
|---|--------------|-----------------|-------------|----------------------|------|---|
| <b>Cancer</b>                           |              |                 |             |                      |      |   |
| Bladder cancer                          | 1601         | 1819            | 10          | 0.62                 | EUR  | NBCS <sup>76</sup>                            |
| Breast cancer                           | 48155        | 43612           | 13          | 1.00                 | EUR  | BCAC <sup>17,77</sup>                         |
| <i>Estrogen receptor –ve</i>            | 7465         | 42175           | 13          | 1.00                 | EUR  | BCAC <sup>17,77</sup>                         |
| <i>Estrogen receptor +ve</i>            | 27074        | 41749           | 13          | 1.00                 | EUR  | BCAC <sup>17,77</sup>                         |
| Colorectal cancer                       | 14537        | 16922           | 9           | 1.00                 | EUR  | CORECT/GECCO <sup>64,78</sup>                 |
| Endometrial cancer                      | 6608         | 37925           | 12          | 1.00                 | EUR  | ECAC <sup>79,80</sup>                         |
| Esophageal SCC                          | 1942         | 2111            | 11          | 0.64                 | EA   | Abnet <sup>81</sup>                           |
| Glioma                                  | 1130         | 6300            | 12          | 0.72                 | EUR  | Wrensch <sup>82</sup> & Walsh <sup>66</sup>   |
| Head & neck cancer                      | 2082         | 3477            | 12          | 1.00                 | EUR  | McKay et al <sup>83</sup>                     |
| Kidney cancer                           | 2461         | 5081            | 12          | 0.99                 | EUR  | KIDRISK <sup>84</sup>                         |
| Lung cancer                             | 11348        | 15861           | 13          | 1.00                 | EUR  | ILCCO <sup>85</sup>                           |
| <i>Adenocarcinoma</i>                   | 3442         | 14894           | 13          | 1.00                 | EUR  | ILCCO <sup>85</sup>                           |
| <i>Squamous cell carcinoma</i>          | 3275         | 15038           | 13          | 1.00                 | EUR  | ILCCO <sup>85</sup>                           |
| Skin cancer                             |              |                 |             |                      |      |   |
| <i>Melanoma</i>                         | 12814        | 23203           | 13          | 1.00                 | EUR  | MC <sup>86</sup>                              |
| <i>Basal cell carcinoma</i>             | 3361         | 11518           | 13          | 1.00                 | EUR  | NHS/HPFS <sup>87</sup>                        |
| Neuroblastoma                           | 2101         | 4202            | 12          | 0.87                 | EUR  | Diskin <sup>88</sup>                          |
| Ovarian cancer                          | 15397        | 30816           | 13          | 1.00                 | EUR  | OCAC <sup>17,89</sup>                         |
| <i>Clear cell</i>                       | 1016         | 30816           | 13          | 0.76                 | EUR  | OCAC <sup>17,89</sup>                         |
| <i>Endometrioid</i>                     | 2154         | 30816           | 13          | 0.98                 | EUR  | OCAC <sup>17,89</sup>                         |
| <i>Mucinous</i>                         | 1643         | 30816           | 13          | 0.94                 | EUR  | OCAC <sup>17,89</sup>                         |
| <i>Serous invasive</i>                  | 9608         | 30816           | 13          | 1.00                 | EUR  | OCAC <sup>17,89</sup>                         |
| <i>Serous LMP</i>                       | 972          | 30816           | 13          | 0.73                 | EUR  | OCAC <sup>17,89</sup>                         |
| Pancreatic cancer                       | 5105         | 8739            | 12          | 1.00                 | EUR  | PanScan (incl. EPIC) <sup>90</sup>            |
| Prostate cancer                         | 22297        | 22323           | 11          | 1.00                 | EUR  | PRACTICAL <sup>91,92</sup>                    |
| Testicular germ cell cancer             | 986          | 4946            | 11          | 0.52                 | EUR  | Turnbull <sup>93</sup> & Rapley <sup>94</sup> |
| <b>Autoimmune/inflammatory diseases</b> |              |                 |             |                      |      |   |
| Alopecia areata                         | 2332         | 5233            | 7           | 0.60                 | EUR  | Betz <sup>95</sup>                            |
| Atopic dermatitis                       | 10788        | 30047           | 13          | 1.00                 | EUR  | EAGLE <sup>96</sup>                           |
| Celiac disease                          | 4533         | 10750           | 3           | 0.82                 | EUR  | Dubois <sup>97</sup>                          |
| Inflammatory bowel disease              |              |                 |             |                      |      |   |
| <i>Crohn's disease</i>                  | 5956         | 14927           | 11          | 1.00                 | EUR  | IIBDGC <sup>98</sup>                          |
| <i>Ulcerative colitis</i>               | 6968         | 20464           | 12          | 1.00                 | EUR  | IIBDGC <sup>98</sup>                          |
| Juvenile idiopathic arthritis           | 1866         | 14786           | 11          | 0.87                 | EUR  | Thompson <sup>99†</sup>                       |
| Multiple sclerosis                      | 14498        | 24091           | 3           | 1.00                 | EUR  | IMSGC <sup>100</sup>                          |
| Aggressive periodontitis                | 888          | 6789            | 13          | 0.63                 | EUR  | Schaefer <sup>101</sup>                       |
| Rheumatoid arthritis                    | 5538         | 20163           | 11          | 1.00                 | EUR  | Stahl <sup>102</sup>                          |
| <b>Cardiovascular diseases</b>          |              |                 |             |                      |      |   |
| Abdominal aortic aneurysm               | 4972         | 99858           | 13          | 1.00                 | EUR  | AC <sup>103–108</sup>                         |
| Coronary heart disease                  | 22233        | 64762           | 13          | 1.00                 | EUR  | CARDIoGRAM <sup>109</sup>                     |
| Heart failure                           | 2526         | 20926           | 13          | 0.99                 | EUR  | CHARGE-HF <sup>110</sup>                      |
| Hemorrhagic stroke                      | 2963         | 5503            | 12          | 0.96                 | EUR  | METASTROKE/ISGC <sup>111</sup>                |
| Ischemic stroke                         | 12389        | 62004           | 13          | 1.00                 | EUR  | METASTROKE/ISGC <sup>112,113</sup>            |
| <i>large vessel disease</i>             | 2167         | 62004           | 13          | 0.99                 | EUR  | METASTROKE/ISGC <sup>112,113</sup>            |
| <i>small vessel disease</i>             | 1894         | 62004           | 13          | 0.97                 | EUR  | METASTROKE/ISGC <sup>112</sup>                |
| <i>cardioembolic</i>                    | 2365         | 62004           | 13          | 0.99                 | EUR  | METASTROKE/ISGC <sup>112</sup>                |
| Sudden cardiac arrest                   | 3954         | 21200           | 13          | 1.00                 | EUR  | Unpublished                                   |
| <b>Diabetes</b>                         |              |                 |             |                      |      |   |
| Type 1 diabetes                         | 7514         | 9045            | 6           | 0.95                 | EUR  | T1DBase <sup>114,115</sup>                    |
| Type 2 diabetes                         | 10415        | 53655           | 11          | 1.00                 | EUR  | DIAGRAM <sup>116</sup>                        |
| <b>Eye disease</b>                      |              |                 |             |                      |      |   |

|  |       |       |    |      |     |                                     |
|--|-------|-------|----|------|-----|-------------------------------------|
| AMD  | 7473  | 51177 | 13 | 1.00 | EUR | AMD Gene <sup>117</sup>             |
| Retinopathy                                | 1122  | 18289 | 12 | 0.75 | EUR | Jensen <sup>118</sup>               |
| <b>Lung diseases</b>                       |       |       |    |      |     |                                     |
| Asthma                                     | 13034 | 20638 | 4  | 1.00 | EUR | Ferreira/GABRIEL <sup>119,120</sup> |
| COPD                                       | 2812  | 2534  | 12 | 0.85 | EUR | COPDGene <sup>121</sup>             |
| Interstitial lung disease                  | 1616  | 4683  | 9  | 0.60 | EUR | Fingerlin <sup>122</sup>            |
| <b>Neurological / psychiatric diseases</b> |       |       |    |      |     |                                     |
| ALS  | 6100  | 7125  | 12 | 1.00 | EUR | SLAGEN/ALSGEN <sup>123</sup>        |
| Alzheimer's disease                        | 17008 | 37154 | 12 | 1.00 | EUR | IGAP <sup>124</sup>                 |
| Anorexia nervosa                           | 2907  | 14860 | 9  | 0.93 | EUR | GCAN <sup>125</sup>                 |
| Autism                                     | 4949  | 5314  | 7  | 0.82 | EUR | PGC <sup>126</sup>                  |
| Bipolar disorder                           | 7481  | 9250  | 9  | 1.00 | EUR | PGC <sup>127</sup>                  |
| Major depressive disorder                  | 9240  | 9519  | 8  | 0.99 | EUR | PGC <sup>128</sup>                  |
| Schizophrenia                              | 35476 | 46839 | 12 | 1.00 | EUR | PGC <sup>129</sup>                  |
| Tourette syndrome                          | 1177  | 4955  | 13 | 0.74 | EUR | TICG/TSAICG <sup>130</sup>          |
| <b>Other</b>                               |       |       |    |      |     |                                     |
| Chronic kidney disease                     | 5807  | 56430 | 13 | 1.00 | EUR | CKDGen <sup>131</sup>               |
| Endometriosis                              | 4604  | 9393  | 11 | 1.00 | Mix | Nyholt <sup>132</sup>               |

---

---

**Study acronyms:** AC, the aneurysm consortium; **ALSGEN**, the International Consortium on Amyotrophic Lateral Sclerosis Genetics; **AMD Gene**, Age-related Macular Degeneration Gene Consortium; **BCAC**, Breast Cancer Association Consortium; **CARDIoGRAM**, Coronary ARtery Disease Genome wide Replication and Meta-analysis; **CHARGE-HF**, Cohorts for Heart and Aging Research in Genomic Epidemiology Consortium – Heart Failure Working Group; **COPDGene**, The Genetic Epidemiology of Chronic Obstructive Pulmonary Disease; **CKDGen**, Chronic Kidney Disease Genetics consortium; **CORECT**, ColoRectal Transdisciplinary Study; **DIAGRAM**, DIAbetes Genetics Replication And Meta-analysis; **EAGLE**, EARly Genetics & Lifecourse Epidemiology Eczema Consortium (excluding 23andMe); **ECAC**, Endometrial Cancer Association Consortium; **EPIC**, European Prospective Investigation into Cancer and Nutrition study; **GABRIEL**, Multidisciplinary Study to Identify the Genetic and Environmental Causes of Asthma in the European Community; **GCAN**, Genetic Consortium for Anorexia Nervosa; **GECCO**, Genetics and Epidemiology of Colorectal Cancer Consortium; **IGAP**, International Genomics of Alzheimer's Project; **HPFS**, Health Professionals Follow-Up Study; **ILCCO**, International Lung Cancer Consortium; **IMSGC**, International Multiple Sclerosis Genetic Consortium; **IIBDGC**, International Inflammatory Bowel Disease Genetics Consortium; **KIDRISK**, Kidney cancer consortium; **MC**, the melanoma meta-analysis consortium; **METASTROKE/ISGC**, METASTROKE project of the International Stroke Genetics Consortium; **NBCS**, Nijmegen Bladder Cancer Study; **NHS**, Nurses' Health Study; **OCAC**, Ovarian Cancer Association Consortium; **PanScan**, Pancreatic Cancer Cohort Consortium; **PGC**, Psychiatric Genomics Consortium; **PRACTICAL**, Prostate Cancer Association Group to Investigate Cancer Associated Alterations in the Genome; **SLAGEN**, Italian Consortium for the Genetics of Ayotrophic Lateral Sclerosis; **T1DBase**, type 1 diabetes database; **TICG** (Tourette International Collaborative-Genetics); **TSAICG** (Tourette Syndrome Association International Consortium for Genetics); **Abbreviations:** **ALS**, amyotrophic lateral sclerosis; **AMD**, age-related macular degeneration; **COPD**, chronic obstructive pulmonary disease; **EUR**, European; **EA**, East Asian; **LMP**, low malignant potential; **No.**, number; **Pop.**, population; **SCC**, squamous cell carcinoma; **SNP**, single nucleotide polymorphism; **-ve**, negative; **+ve**, positive; †plus previously unpublished data.

## Figure 1. The association between genetically increased telomere length and odds of primary non-communicable diseases

### Legend to Figure 1

\*P value for association between genetically increased telomere length and disease from maximum likelihood; the effect estimate for heart failure is a hazard ratio (all others are odds ratios);  $P_{het}$ , P-value for heterogeneity amongst SNPs within the instrument; COPD, chronic obstructive pulmonary disease; SNP, single nucleotide polymorphism; CI, confidence interval; LMP, low malignancy potential; ER, estrogen receptor; -VE, negative; +VE, positive.

## Figure 2. The association between genetically increased telomere length and odds of cancer as a function of selected characteristics

### Legend to Figure 2

The plotted data show how the strength of the relationship between genetically increased telomere length and cancer varies by the selected characteristic. The  $R^2$  statistic indicates how much of the variation between cancers can be explained by the selected characteristic. P-values are from meta-regression models. Circle sizes are proportional to the inverse of the variance of the log odds ratio. The hashed line indicates the null of no association between telomere length and cancer (i.e. an odds ratio of 1). Data for percentage survival 5 years after diagnosis, cancer incidence and median age-at-diagnosis was downloaded from the Surveillance, Epidemiology, and End Results Program.<sup>33</sup> Data for average lifetime number of stem cell divisions was downloaded from Tomasetti and Vogelstein.<sup>34</sup> Not all cancers had information available for the selected characteristics (hence the number of cancers varies across the subplots). Information was available for 9 cancers for tissue-specific rates of stem cell division, 13 cancers for percentage surviving 5 years post-diagnosis, 17 cancers for cancer incidence and 13 cancers for median age-at-diagnosis. SD, standard deviation; OR, Odds ratio.

## Figure 3. Comparison of genetic and prospective observational studies<sup>†</sup> of the association between telomere length and disease

### Legend to Figure 3

\*from fixed-effects meta-analysis of independent observational studies described in Table S3; <sup>†</sup>search strategy and characteristics for observational studies are described in Tables S3 and S4; ‡CCHS and CGPS; +PLCO, ATBC & SWHS (acronyms explained in Table S3); CI, confidence interval

## REFERENCES

1. Blackburn EH, Epel ES, Lin J. Human telomere biology: A contributory and interactive factor in aging, disease risks, and protection. *Science* (80- ) 2015;350(6265):1193–8.
2. López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging. *Cell* 2013;153(6):1194–217.
3. Samani NJ, van der Harst P. Biological ageing and cardiovascular disease. *Heart* 2008;94(5):537–9.
4. Weischer M, Bojesen SE, Nordestgaard BG. Telomere shortening unrelated to smoking, body weight, physical activity, and alcohol intake: 4,576 general population individuals with repeat measurements 10 years apart. *PLoS Genet* 2014;10(3):e1004191.
5. Houben MJ, Moonen HJJ, van Schooten FJ, Hageman GJ. Telomere length assessment: biomarker of chronic oxidative stress? *Free Radic Biol Med* 2008;44(3):235–46.
6. Marchesi V. Risk factors: Short telomeres: association with cancer survival and risk. *Nat Rev Clin Oncol* 2013;10(5):247.
7. Haycock PC, Heydon EE, Kaptoge S, Butterworth a. S, Thompson A, Willeit P. Leucocyte telomere length and risk of cardiovascular disease: systematic review and meta-analysis. *BMJ* 2014;349(jul08 3):g4227–g4227.
8. Rode L, Nordestgaard BG, Bojesen SE. Peripheral blood leukocyte telomere length and mortality among 64,637 individuals from the general population. *J Natl Cancer Inst* 2015;107(6):djv074.
9. Zhao J, Miao K, Wang H, Ding H, Wang DW. Association between telomere length and type 2 diabetes mellitus: a meta-analysis. *PLoS One* 2013;8(11):e79993.
10. Weischer M, Nordestgaard BG, Cawthon RM, Freiberg JJ, Tybjaerg-Hansen A, Bojesen SE. Short telomere length, cancer survival, and cancer risk in 47102 individuals. *J Natl Cancer Inst* 2013;105(7):459–68.
11. Ma H, Zhou Z, Wei S, et al. Shortened telomere length is associated with increased risk of cancer: a meta-analysis. *PLoS One* 2011;6(6):e20466.
12. Wentzensen IM, Mirabello L, Pfeiffer RM, Savage SA. The association of telomere length and cancer: a meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2011;20(6):1238–50.
13. Pooley KA, Sandhu MS, Tyrer J, et al. Telomere length in prospective and retrospective cancer case-control studies. *Cancer Res* 2010;70(8):3170–6.
14. Hou L, Joyce BT, Gao T, et al. Blood Telomere Length Attrition and Cancer Development in the Normative Aging Study Cohort. *EBioMedicine* 2015;2(6):591–6.
15. Welter D, MacArthur J, Morales J, et al. The NHGRI GWAS Catalog, a curated resource of SNP-trait associations. *Nucleic Acids Res* 2014;42(Database issue):D1001–6.

- 494 16. Burdett T, Hall P, Hastings E, et al. The NHGRI-EBI Catalog of published genome-  
495 wide association studies [Internet]. [cited 2015 Jan 15];Available from:  
496 [www.ebi.ac.uk/gwas](http://www.ebi.ac.uk/gwas)
- 497 17. Pooley KA, Bojesen SE, Weischer M, et al. A genome-wide association scan (GWAS)  
498 for mean telomere length within the COGS project: identified loci show little  
499 association with hormone-related cancer risk. *Hum Mol Genet* 2013;22(24):5056–64.
- 500 18. Mangino M, Hwang S-J, Spector TD, et al. Genome-wide meta-analysis points to  
501 CTC1 and ZNF676 as genes regulating telomere homeostasis in humans. *Hum Mol*  
502 *Genet* 2012;21(24):5385–94.
- 503 19. Prescott J, Kraft P, Chasman DI, et al. Genome-wide association study of relative  
504 telomere length. *PLoS One* 2011;6(5):e19635.
- 505 20. Gu J, Chen M, Shete S, et al. A genome-wide association study identifies a locus on  
506 chromosome 14q21 as a predictor of leukocyte telomere length and as a marker of  
507 susceptibility for bladder cancer. *Cancer Prev Res (Phila)* 2011;4(4):514–21.
- 508 21. Codd V, Nelson CP, Albrecht E, et al. Identification of seven loci affecting mean  
509 telomere length and their association with disease. *Nat Genet* 2013;45(4):422–7.
- 510 22. Codd V, Mangino M, van der Harst P, et al. Common variants near TERC are  
511 associated with mean telomere length. *Nat Genet* 2010;42(3):197–9.
- 512 23. Liu Y, Cao L, Li Z, et al. A genome-wide association study identifies a locus on TERT  
513 for mean telomere length in Han Chinese. *PLoS One* 2014;9(1):e85043.
- 514 24. Saxena R, Bjorndal A, Prescott J, et al. Genome-wide association study identifies  
515 variants in casein kinase II (CSNK2A2) to be associated with leukocyte telomere  
516 length in a Punjabi Sikh diabetic cohort. *Circ Cardiovasc Genet* 2014;7(3):287–95.
- 517 25. Levy D, Neuhausen SL, Hunt SC, et al. Genome-wide association identifies OBFC1 as  
518 a locus involved in human leukocyte telomere biology. *Proc Natl Acad Sci U S A*  
519 2010;107(20):9293–8.
- 520 26. Hindorff LA, MacArthur J, Morales J, et al. A catalog of published genome-wide  
521 association studies [Internet]. [cited 2015 Jan 15];Available from:  
522 [www.genome.gov/gwastudies](http://www.genome.gov/gwastudies)
- 523 27. Chene G, Thompson SG. Methods for Summarizing the Risk Associations of  
524 Quantitative Variables in Epidemiologic Studies in a Consistent Form. *Am J*  
525 *Epidemiol* 1996;144(6):610–21.
- 526 28. Burgess S, Scott RA, Timpson NJ, Davey Smith G, Thompson SG, EPIC-InterAct  
527 Consortium. Using published data in Mendelian randomization: a blueprint for  
528 efficient identification of causal risk factors. *Eur J Epidemiol* 2015;30(7):543–52.
- 529 29. Thomas DC, Lawlor DA, Thompson JR. Re: Estimation of bias in nongenetic  
530 observational studies using “Mendelian triangulation” by Bautista et al. *Ann*  
531 *Epidemiol* 2007;17(7):511–3.
- 532 30. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid  
533 instruments: effect estimation and bias detection through Egger regression. *Int J*  
534 *Epidemiol* 2015;44(2):512–25.



- 535 31. VanderWeele TJ, Tchetgen Tchetgen EJ, Cornelis M, Kraft P. Methodological  
536 challenges in mendelian randomization. *Epidemiology* 2014;25(3):427–35.
- 537 32. Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent estimation in  
538 Mendelian randomization with some invalid instruments using a weighted median  
539 estimator. *Genet Epidemiol*
- 540 33. National Cancer Institute. Surveillance, Epidemiology, and End Results Program  
541 [Internet]. [cited 2015 Aug 1];Available from: [www.seer.cancer.gov](http://www.seer.cancer.gov)
- 542 34. Tomasetti C, Vogelstein B. Variation in cancer risk among tissues can be explained by  
543 the number of stem cell divisions. *Science* 2015;347(6217):78–81.
- 544 35. U.S. Census Bureau [Internet]. [cited 2016 Jul 11];Available from: U.S. Census  
545 Bureau
- 546 36. R Core Team. A language and environment for statistical computing. 2013;
- 547 37. Burgess S, Thompson SG. Bias in causal estimates from Mendelian randomization  
548 studies with weak instruments. *Stat Med* 2011;30(11):1312–23.
- 549 38. Davey Smith G, Ebrahim S. What can mendelian randomisation tell us about  
550 modifiable behavioural and environmental exposures? *BMJ Br Med J*  
551 2005;330(7499):1076–9.
- 552 39. Anic GM, Sondak VK, Messina JL, et al. Telomere length and risk of melanoma,  
553 squamous cell carcinoma, and basal cell carcinoma. *Cancer Epidemiol*  
554 2013;37(4):434–9.
- 555 40. Pellatt AJ, Wolff RK, Torres-Mejia G, et al. Telomere length, telomere-related genes,  
556 and breast cancer risk: the breast cancer health disparities study. *Genes Chromosomes*  
557 *Cancer* 2013;52(7):595–609.
- 558 41. Caini S, Raimondi S, Johansson H, et al. Telomere length and the risk of cutaneous  
559 melanoma and non-melanoma skin cancer: a review of the literature and meta-analysis.  
560 *J Dermatol Sci* 2015;80(3):168–74.
- 561 42. Sanchez-Espiridion B, Chen M, Chang JY, et al. Telomere length in peripheral blood  
562 leukocytes and lung cancer risk: a large case-control study in Caucasians. *Cancer Res*  
563 2014;74(9):2476–86.
- 564 43. Campa D, Mergarten B, De Vivo I, et al. Leukocyte telomere length in relation to  
565 pancreatic cancer risk: a prospective study. *Cancer Epidemiol Biomarkers Prev*  
566 2014;23(11):2447–54.
- 567 44. Cui Y, Cai Q, Qu S, et al. Association of leukocyte telomere length with colorectal  
568 cancer risk: nested case-control findings from the Shanghai Women’s Health Study.  
569 *Cancer Epidemiol Biomarkers Prev* 2012;21(10):1807–13.
- 570 45. De Vivo I, Prescott J, Wong JYY, Kraft P, Hankinson SE, Hunter DJ. A prospective  
571 study of relative telomere length and postmenopausal breast cancer risk. *Cancer*  
572 *Epidemiol Biomarkers Prev* 2009;18(4):1152–6.
- 573 46. Han J, Qureshi AA, Prescott J, et al. A prospective study of telomere length and the  
574 risk of skin cancer. *J Invest Dermatol* 2009;129(2):415–21.
- 575 47. Hofmann JN, Lan Q, Cawthon R, et al. A prospective study of leukocyte telomere

length and risk of renal cell carcinoma. *Cancer Epidemiol Biomarkers Prev* 2013;22(5):997–1000.

48. Julin B, Shui I, Heaphy CM, et al. Circulating leukocyte telomere length and risk of overall and aggressive prostate cancer. *Br J Cancer* 2015;112(4):769–76.

49. Kim S, Sandler DP, Carswell G, et al. Telomere length in peripheral blood and breast cancer risk in a prospective case-cohort analysis: results from the Sister Study. *Cancer Causes Control* 2011;22(7):1061–6.

50. Lan Q, Cawthon R, Shen M, et al. A prospective study of telomere length measured by monochrome multiplex quantitative PCR and risk of non-Hodgkin lymphoma. *Clin Cancer Res* 2009;15(23):7429–33.

51. Lee I-M, Lin J, Castonguay AJ, Barton NS, Buring JE, Zee RYL. Mean leukocyte telomere length and risk of incident colorectal carcinoma in women: a prospective, nested case-control study. *Clin Chem Lab Med* 2010;48(2):259–62.

52. Liang G, Qureshi AA, Guo Q, De Vivo I, Han J. No association between telomere length in peripheral blood leukocytes and the risk of nonmelanoma skin cancer. *Cancer Epidemiol Biomarkers Prev* 2011;20(5):1043–5.

53. Lynch SM, Major JM, Cawthon R, et al. A prospective analysis of telomere length and pancreatic cancer in the alpha-tocopherol beta-carotene cancer (ATBC) prevention study. *Int J Cancer* 2013;133(11):2672–80.

54. McGrath M, Wong JYY, Michaud D, Hunter DJ, De Vivo I. Telomere length, cigarette smoking, and bladder cancer risk in men and women. *Cancer Epidemiol Biomarkers Prev* 2007;16(4):815–9.

55. Nan H, Du M, De Vivo I, et al. Shorter telomeres associate with a reduced risk of melanoma development. *Cancer Res* 2011;71(21):6758–63.

56. Prescott J, McGrath M, Lee I-M, Buring JE, De Vivo I. Telomere length and genetic analyses in population-based studies of endometrial cancer risk. *Cancer* 2010;116(18):4275–82.

57. Qu S, Wen W, Shu X-O, et al. Association of leukocyte telomere length with breast cancer risk: nested case-control findings from the Shanghai Women’s Health Study. *Am J Epidemiol* 2013;177(7):617–24.

58. Risques RA, Vaughan TL, Li X, et al. Leukocyte telomere length predicts cancer risk in Barrett’s esophagus. *Cancer Epidemiol Biomarkers Prev* 2007;16(12):2649–55.

59. Seow WJ, Cawthon RM, Purdue MP, et al. Telomere length in white blood cell DNA and lung cancer: a pooled analysis of three prospective cohorts. *Cancer Res* 2014;74(15):4090–8.

60. Shen M, Cawthon R, Rothman N, et al. A prospective study of telomere length measured by monochrome multiplex quantitative PCR and risk of lung cancer. *Lung Cancer* 2011;73(2):133–7.

61. Walcott F, Rajaraman P, Gadalla SM, et al. Telomere length and risk of glioma. *Cancer Epidemiol* 2013;37(6):935–8.

62. Zee RYL, Castonguay AJ, Barton NS, Buring JE. Mean telomere length and risk of

617 incident colorectal carcinoma: a prospective, nested case-control approach. *Cancer*  
618 *Epidemiol Biomarkers Prev* 2009;18(8):2280–2.

619 63. Willeit P, Willeit J, Mayr A, et al. Telomere length and risk of incident cancer and  
620 cancer mortality. *Jama* 2010;304(1538–3598 (Electronic)):69–75.

621 64. Zhang C, Doherty J a., Burgess S, et al. Genetic determinants of telomere length and  
622 risk of common cancers: a Mendelian randomization study. *Hum Mol Genet*  
623 2015;24(18):5356–66.

624 65. Iles MM, Bishop DT, Taylor JC, et al. The effect on melanoma risk of genes  
625 previously associated with telomere length. *J Natl Cancer Inst* 2014;106(10).

626 66. Walsh KM, Codd V, Rice T, et al. Longer genotypically-estimated leukocyte telomere  
627 length is associated with increased adult glioma risk. *Oncotarget* 2015;6(40):42468–  
628 77.

629 67. Ojha J, Codd V, Nelson CP, et al. Genetic Variation Associated with Longer Telomere  
630 Length Increases Risk of Chronic Lymphocytic Leukemia. *Cancer Epidemiol*  
631 *Biomarkers Prev* 2016;25(7):1043–9.

632 68. Armanios M, Blackburn EH. The telomere syndromes. *Nat Rev Genet*  
633 2012;13(10):693–704.

634 69. Armanios M. Syndromes of telomere shortening. *Annu Rev Genomics Hum Genet*  
635 2009;10(46):45–61.

636 70. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell*  
637 2011;144(5):646–74.

638 71. Blasco MA. Telomere length, stem cells and aging. *Nat Chem Biol* 2007;3(10):640–9.

639 72. Stone RC, Horvath K, Kark JD, Susser E, Tishkoff SA, Aviv A. Telomere Length and  
640 the Cancer-Atherosclerosis Trade-Off. *PLoS Genet* 2016;12(7):e1006144.

641 73. Pierce BL, Burgess S. Efficient design for Mendelian randomization studies:  
642 subsample and 2-sample instrumental variable estimators. *Am J Epidemiol*  
643 2013;178(7):1177–84.

644 74. Burgess S, Butterworth AS, Thompson JR. Beyond Mendelian randomization: how to  
645 interpret evidence of shared genetic predictors. *J Clin Epidemiol* 2015;1–9.

646 75. Centers for Disease Control and Prevention [Internet]. [cited 2016 Jul 14];Available  
647 from: <http://www.cdc.gov/nchs/fastats/deaths.htm>

648 76. Rafnar T, Sulem P, Thorleifsson G, et al. Genome-wide association study yields  
649 variants at 20p12.2 that associate with urinary bladder cancer. *Hum Mol Genet*  
650 2014;23(20):5545–57.

651 77. Michailidou K, Hall P, Gonzalez-Neira A, et al. Large-scale genotyping identifies 41  
652 new loci associated with breast cancer risk. *Nat Genet* 2013;45(4):353–61, 361–2.

653 78. Schumacher FR, Schmit SL, Jiao S, et al. Genome-wide association study of colorectal  
654 cancer identifies six new susceptibility loci. *Nat Commun* 2015;6:7138.

655 79. Spurdle AB, Thompson DJ, Ahmed S, et al. Genome-wide association study identifies  
656 a common variant associated with risk of endometrial cancer. *Nat Genet*

2011;43(5):451–4.

80. Painter JN, O’Mara TA, Batra J, et al. Fine-mapping of the HNF1B multicancer locus identifies candidate variants that mediate endometrial cancer risk. *Hum Mol Genet* 2015;24(5):1478–92.

81. Abnet CC, Freedman ND, Hu N, et al. A shared susceptibility locus in PLCE1 at 10q23 for gastric adenocarcinoma and esophageal squamous cell carcinoma. *Nat Genet* 2010;42(9):764–7.

82. Wrensch M, Jenkins RB, Chang JS, et al. Variants in the CDKN2B and RTEL1 regions are associated with high-grade glioma susceptibility. *Nat Genet* 2009;41(8):905–8.

83. McKay JD, Truong T, Gaborieau V, et al. A genome-wide association study of upper aerodigestive tract cancers conducted within the INHANCE consortium. *PLoS Genet* 2011;7(3):e1001333.

84. Purdue MP, Johansson M, Zelenika D, et al. Genome-wide association study of renal cell carcinoma identifies two susceptibility loci on 2p21 and 11q13.3. *Nat Genet* 2010;43(1):60–5.

85. Wang Y, McKay JD, Rafnar T, et al. Rare variants of large effect in BRCA2 and CHEK2 affect risk of lung cancer. *Nat Genet* 2014;46(7).

86. Law MH, Bishop DT, Lee JE, et al. Genome-wide meta-analysis identifies five new susceptibility loci for cutaneous malignant melanoma. *Nat Genet* 2015;47(9):987–95.

87. Zhang M, Song F, Liang L, et al. Genome-wide association studies identify several new loci associated with pigmentation traits and skin cancer risk in European Americans. *Hum Mol Genet* 2013;22(14):2948–59.

88. Diskin SJ, Capasso M, Schnepf RW, et al. Common variation at 6q16 within HACE1 and LIN28B influences susceptibility to neuroblastoma. *Nat Genet* 2012;44(10):1126–30.

89. Pharoah PDP, Tsai Y-Y, Ramus SJ, et al. GWAS meta-analysis and replication identifies three new susceptibility loci for ovarian cancer. *Nat Genet* 2013;45(4):362–70, 370-2.

90. Wolpin BM, Rizzato C, Kraft P, et al. Genome-wide association study identifies multiple susceptibility loci for pancreatic cancer. *Nat Genet* 2014;46(9):994–1000.

91. Eeles RA, Olama AA Al, Benlloch S, et al. Identification of 23 new prostate cancer susceptibility loci using the iCOGS custom genotyping array. *Nat Genet* 2013;45(4):385–91, 391-2.

92. Al Olama AA, Kote-Jarai Z, Berndt SI, et al. A meta-analysis of 87,040 individuals identifies 23 new susceptibility loci for prostate cancer. *Nat Genet* 2014;46(10):1103–9.

93. Turnbull C, Rapley E a, Seal S, et al. Variants near DMRT1, TERT and ATF7IP are associated with testicular germ cell cancer. *Nat Genet* 2010;42(7):604–7.

94. Rapley EA, Turnbull C, Al Olama AA, et al. A genome-wide association study of testicular germ cell tumor. *Nat Genet* 2009;41(7):807–10.

95. Betz RC, Petukhova L, Ripke S, et al. Genome-wide meta-analysis in alopecia areata resolves HLA associations and reveals two new susceptibility loci. *Nat Commun* 2015;6:5966.
96. EARly Genetics and Lifecourse Epidemiology (EAGLE) Eczema Consortium, Australian Asthma Genetics Consortium(AAGC), Australian Asthma Genetics Consortium AAGC. Multi-ancestry genome-wide association study of 21,000 cases and 95,000 controls identifies new risk loci for atopic dermatitis. *Nat Genet* 2015;47(12):1449–56.
97. Dubois PC a, Trynka G, Franke L, et al. Multiple common variants for celiac disease influencing immune gene expression. *Nat Genet* 2010;42(4):295–302.
98. Liu JZ, van Sommeren S, Huang H, et al. Association analyses identify 38 susceptibility loci for inflammatory bowel disease and highlight shared genetic risk across populations. *Nat Genet* 2015;47(9):979–86.
99. Thompson SD, Marion MC, Sudman M, et al. Genome-wide association analysis of juvenile idiopathic arthritis identifies a new susceptibility locus at chromosomal region 3q13. *Arthritis Rheum* 2012;64(8):2781–91.
100. Beecham AH, Patsopoulos NA, Xifara DK, et al. Analysis of immune-related loci identifies 48 new susceptibility variants for multiple sclerosis. *Nat Genet* 2013;45(11):1353–60.
101. Schaefer AS, Richter GM, Nothnagel M, et al. A genome-wide association study identifies GLT6D1 as a susceptibility locus for periodontitis. *Hum Mol Genet* 2010;19(3):553–62.
102. Stahl EA, Raychaudhuri S, Remmers EF, et al. Genome-wide association study meta-analysis identifies seven new rheumatoid arthritis risk loci. *Nat Genet* 2010;42(6):508–14.
103. Bown MJ, Jones GT, Harrison SC, et al. Abdominal aortic aneurysm is associated with a variant in low-density lipoprotein receptor-related protein 1. *Am J Hum Genet* 2011;89(5):619–27.
104. Gretarsdottir S, Baas AF, Thorleifsson G, et al. Genome-wide association study identifies a sequence variant within the DAB2IP gene conferring susceptibility to abdominal aortic aneurysm. *Nat Genet* 2010;42(8):692–7.
105. Jones GT, Bown MJ, Gretarsdottir S, et al. A sequence variant associated with sortilin-1 (SORT1) on 1p13.3 is independently associated with abdominal aortic aneurysm. *Hum Mol Genet* 2013;22(14):2941–7.
106. Harrison SC, Smith AJP, Jones GT, et al. Interleukin-6 receptor pathways in abdominal aortic aneurysm. *Eur Heart J* 2013;34(48):3707–16.
107. Elmore JR, Obmann MA, Kuivaniemi H, et al. Identification of a genetic variant associated with abdominal aortic aneurysms on chromosome 3p12.3 by genome wide association. *J Vasc Surg* 2009;49(6):1525–31.
108. Borthwick K, Smelser D, Bock J, et al. Ephenotyping for Abdominal Aortic Aneurysm in the Electronic Medical Records and Genomics (eMERGE) Network: Algorithm Development and Konstanz Information Miner Workflow. *Int J Biomed Data Min* 2015;4(1).

109. Schunkert H, König IR, Kathiresan S, et al. Large-scale association analysis identifies 13 new susceptibility loci for coronary artery disease. *Nat Genet* 2011;43(4):333–8.
110. Smith NL, Felix JF, Morrison AC, et al. Association of genome-wide variation with the risk of incident heart failure in adults of European and African ancestry: a prospective meta-analysis from the cohorts for heart and aging research in genomic epidemiology (CHARGE) consortium. *Circ Cardiovasc Genet* 2010;3(3):256–66.
111. Woo D, Falcone GJ, Devan WJ, et al. Meta-analysis of genome-wide association studies identifies 1q22 as a susceptibility locus for intracerebral hemorrhage. *Am J Hum Genet* 2014;94(4):511–21.
112. Malik R, Freilinger T, Winsvold BS, et al. Shared genetic basis for migraine and ischemic stroke: A genome-wide analysis of common variants. *Neurology* 2015;84(21):2132–45.
113. Traylor M, Farrall M, Holliday EG, et al. Genetic risk factors for ischaemic stroke and its subtypes (the METASTROKE collaboration): a meta-analysis of genome-wide association studies. *Lancet Neurol* 2012;11(11):951–62.
114. Barrett JC, Clayton DG, Concannon P, et al. Genome-wide association study and meta-analysis find that over 40 loci affect risk of type 1 diabetes. *Nat Genet* 2009;41(6):703–7.
115. Burren OS, Adlem EC, Achuthan P, Christensen M, Coulson RMR, Todd JA. T1DBase: update 2011, organization and presentation of large-scale data sets for type 1 diabetes research. *Nucleic Acids Res* 2011;39(Database):D997–1001.
116. Morris ADPDPA, Voight BFB, Teslovich TMT, et al. Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes. *Nat Genet* 2012;44(9):981–90.
117. Fritsche LG, Chen W, Schu M, et al. Seven new loci associated with age-related macular degeneration. *Nat Genet* 2013;45(4):433–9, 439–2.
118. Jensen RA, Sim X, Li X, et al. Genome-wide association study of retinopathy in individuals without diabetes. *PLoS One* 2013;8(2):e54232.
119. Moffatt MF, Kabesch M, Liang L, et al. Genetic variants regulating ORMDL3 expression contribute to the risk of childhood asthma. *Nature* 2007;448(7152):470–3.
120. Ferreira MAR, Matheson MC, Duffy DL, et al. Identification of IL6R and chromosome 11q13.5 as risk loci for asthma. *Lancet (London, England)* 2011;378(9795):1006–14.
121. Cho MH, McDonald M-LN, Zhou X, et al. Risk loci for chronic obstructive pulmonary disease: a genome-wide association study and meta-analysis. *Lancet Respir Med* 2014;2(3):214–25.
122. Fingerlin TE, Murphy E, Zhang W, et al. Genome-wide association study identifies multiple susceptibility loci for pulmonary fibrosis. *Nat Genet* 2013;45(6):613–20.
123. Fogh I, Ratti A, Gellera C, et al. A genome-wide association meta-analysis identifies a novel locus at 17q11.2 associated with sporadic amyotrophic lateral sclerosis. *Hum Mol Genet* 2014;23(8):2220–31.

782 124. Lambert JC, Ibrahim-Verbaas CA, Harold D, et al. Meta-analysis of 74,046 individuals  
783 identifies 11 new susceptibility loci for Alzheimer's disease. *Nat Genet*  
784 2013;45(12):1452–8.

785 125. Boraska V, Franklin CS, Floyd JAB, et al. A genome-wide association study of  
786 anorexia nervosa. *Mol Psychiatry* 2014;19(10):1085–94.

787 126. Smoller JW, Hospital MG. Identification of risk loci with shared effects on five major  
788 psychiatric disorders: a genome-wide analysis. *Lancet* 2013;381(9875):1371–9.

789 127. Sklar P, Ripke S, Scott LJ, et al. Large-scale genome-wide association analysis of  
790 bipolar disorder identifies a new susceptibility locus near ODZ4. *Nat Genet*  
791 2011;43(10):977–83.

792 128. Ripke S, Wray NR, Lewis CM, et al. A mega-analysis of genome-wide association  
793 studies for major depressive disorder. *Mol Psychiatry* 2013;18(4):497–511.

794 129. Ripke S, Neale BM, Corvin A, et al. Biological insights from 108 schizophrenia-  
795 associated genetic loci. *Nature* 2014;511(7510):421–7.

796 130. Scharf JM, Yu D, Mathews CA, et al. Genome-wide association study of Tourette's  
797 syndrome. *Mol Psychiatry* 2013;18(6):721–8.

798 131. Köttgen A, Pattaro C, Böger C a, et al. New loci associated with kidney function and  
799 chronic kidney disease. *Nat Genet* 2010;42(5):376–84.

800 132. Nyholt DR, Low S-K, Anderson CA, et al. Genome-wide association meta-analysis  
801 identifies new endometriosis risk loci. *Nat Genet* 2012;44(12):1355–9.

802







